

C-Groups^{*}

Kazem Mahdavi[†], Margrethe Flanders[‡], Avi Silterra[§], Mihai Tohaneanu[¶]

February 1, 2008

Abstract

In this paper we explore the structure and properties of C-groups. We define a C-group as a group G with $rk(G) < rk(Z(G))$. Using GAP (a group theory program) and traditional methods, we identified an interesting infinite class of C-groups. In particular, we have proved that there is always a C-group of order an integer multiple of a fifth power of a prime. One way to obtain C-groups is take the direct product of certain C-groups, mentioned in this paper, with other appropriate groups. We, at this time, do not know whether the C-groups discussed in this paper are the building block of all finite C-groups. But a complete classification of finite or infinite C-groups is an interesting problem. We have also formulated a number of open questions relating to C-groups: Are they all solvable? What is the structure of the C-groups that are not in our class? Is the minimal number of generators of the center always polynomially bounded by the minimal number of generators of the group? What are the isoperimetric inequalities of infinite C-groups?

Introduction

Call the minimum cardinality of a generating set for a group G the rank of G and denote it by $rk(G)$.

In this paper we study a class of finitely generated groups G such that $rk(G) < rk(Z(G))$ where $Z(G)$ is the center of G . We call these groups **C-groups**.

We came across this class of groups while trying to study finitely generated groups with infinitely generated centers. One class of such groups is the set of 4×4 upper-triangular matrices over the set of rationals with a power of a prime

^{*}Research funded by grants from the NSF grant number NSF(DMS-0097113), and NSA, grant number NSA(MSPR-01IC-180).

[†]Project Mentor, mahdavi@potsdam.edu

[‡]Group Leader, mebflanders@att.net

[§]ajs2@andrew.cmu.edu

[¶]mihai@dartmouth.edu

p in the denominator:

$$\begin{pmatrix} 1 & * & * & * \\ 0 & u_1 & * & * \\ 0 & 0 & u_2 & * \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

where u_i denotes a positive unit. (For further details, see [1]).

We used GAP, a system for computational group theory, both to find finite C-groups of order less than 1000 (with the exceptions of 512 and 768) and to gain insight into their structures [3]. In this paper we explain the results of these investigations.

1 Definitions and Notation

Definition 1 (Elementary abelian). A group G is elementary abelian if $G = \prod_{i < n} \mathbb{Z}_p$ for some $n \in \mathbb{N}$ and p a prime.

Definition 2 (Frattini Subgroup). The **Frattini subgroup** of G , denoted $\Phi(G)$, is the intersection of all maximal subgroups of G when they exist and G otherwise.

Definition 3 (Nilpotent). Let G be a group, define $Z_1(G) = Z(G)$ and $Z_i(G)$ is the inverse image of $Z(G/Z_{i-1}(G))$ under the canonical projection $G \rightarrow G/Z_{i-1}(G)$. The group G is **nilpotent** (of class n) if $Z_n(G) = G$ for some n .

Definition 4 (Commutator). Let G be a group and let $x, y \in G$. Then $[x, y] = xyx^{-1}y^{-1}$ is called the **commutator** of x and y . The subgroup of G generated by the set $\{[x, y] \mid x, y \in G\}$ is called the **commutator subgroup** of G and will be denoted G' .

Definition 5 (Solvable). Let G be a group, and let $G^{(1)} := G'$. Then for $i \geq 1$, define $G^{(i)}$ by $G^{(i)} = (G^{(i-1)})'$. The group G is said to be **solvable** if $G^{(n)} = \langle e \rangle$ for some n .

Definition 6 (p -Group). A group in which every element has order a power of some fixed prime p is called a **p -group**.

Lemma 1. The set $\mathbb{Z}_{n_1} \times \mathbb{Z}_{n_2} \times \dots \times \mathbb{Z}_{n_k}$ with $\gcd(n_1, n_2, \dots, n_k) > 1$ has a minimum of k generators.

Lemma 2. Any group of order p^2 for p a prime is cyclic or elementary abelian.

Theorem 1. A finite group G is a p -group $\Leftrightarrow$ the order of G is a power of some prime p .

Theorem 2. Every finite p -group is nilpotent.

For more information on nilpotent and solvable groups, see [4], pages 100-107.

Theorem 3. Burnside's Basis Theorem *If G is a finite p -group, then $\forall g \in G, g^p \in \Phi(G)$, $G' \leq \Phi(G)$, $\bar{G} = G/\Phi(G)$ is elementary abelian, and $rk(\bar{G}) = rk(G)$.*

Universal Mapping Property of Frattini Subgroup *If H is a finite p -group, $N \triangleleft H$ and H/N is elementary abelian then $\Phi(H) \leq N$.*

See [2] or [5] for details.

2 Main Results

Theorem 4 (Main Result). *Let $G = \mathbb{Z}_{n_1} \times \mathbb{Z}_{n_2} \times \mathbb{Z}_{n_3}$ where $n_1 \mid n_2, n_1 \mid n_3$, and $\gcd(n_1, \frac{n_2}{n_1}, \frac{n_3}{n_1}) > 1$, and binary operation*

$$(x_1, y_1, z_1) \cdot (x_2, y_2, z_2) = ((x_1 + x_2 + y_2 z_1) \bmod n_1, (y_1 + y_2) \bmod n_2, (z_1 + z_2) \bmod n_3).$$

Then $(G, \cdot)$ is a C-group.

Proof. It is easy to verify that G is a group under the above binary operation where $(0, 0, 0)$ is the identity element and

$$(x, y, z)^{-1} = ((n_1 - x) + yz, n_2 - y, n_3 - z).$$

The generators of G are $a = (0, 1, 0)$ and $b = (0, 0, 1)$, since $(1, 0, 0) = bab^{-1}a^{-1}$, and they satisfy the following relations:

$$a^{n_2} = b^{n_3} = (0, 0, 0), (bab^{-1}a^{-1})^{n_1} = (0, 0, 0), a^{n_1}b = ba^{n_1}, ab^{n_1} = b^{n_1}a.$$

Therefore, the canonical form of words in G is

$$a^{k_1 n_1} b^{k_2 n_1} [a, b]^{k_3} a^{k_4} b^{k_5}$$

where $k_1 = 0, 1, \dots, \frac{n_2}{n_1}$, $k_2 = 0, 1, \dots, \frac{n_3}{n_1}$, and $k_3, k_4, k_5 = 0, 1, \dots, n_1 - 1$. Thus the order of G is $n_1 n_2 n_3$.

Let $C = Z(G)$. Then

$$C = \{(x, n_1 y, n_1 z) \mid x \in \mathbb{Z}_{n_1}, y \in \mathbb{Z}_{n_2}, z \in \mathbb{Z}_{n_3}\}.$$

So, $C \cong \mathbb{Z}_{n_1} \times \mathbb{Z}_{\frac{n_2}{n_1}} \times \mathbb{Z}_{\frac{n_3}{n_1}}$ of order $n_1(\frac{n_2}{n_1})(\frac{n_3}{n_1})$ by Lemma 1.

Therefore, G is a C-group of order $n_1 n_2 n_3$. □

Remark. This defines an infinite class of C-groups which we call α C-groups.

Theorem 5. *There is a unique C-group of order p^5 for any prime p .*

Proof. Existence follows from Theorem 4 with $n_1 = p, n_2 = p^2, n_3 = p^2$.

Uniqueness is more difficult. Suppose G is a C-group such that $|G| = p^5$. First we'll show $rk(Z(G)) = 3$ and $rk(G) = 2$. Let $|G/Z(G)| = p^q$. $q = 1$ implies G abelian, and $q \geq 3$ implies that $rk(Z(G)) \leq 2$. Hence, $q = 2$ and

so $|Z(G)| = p^3$. Therefore, $rk(Z(G)) \leq 3$, so as G is a C-group we have $rk(Z(G)) = 3$ and $rk(G) = 2$.

Next we will show that $\Phi(G) = Z(G)$, though we'll only need $\Phi(G) \leq Z(G)$. To begin, note that $G/Z(G)$ is elementary abelian by Lemma 2. By the UMP of the Frattini subgroup, we have $\Phi(G) \leq Z(G)$. Next by the Burnside's Basis Theorem and $rk(G) = 2$, we must have $|\Phi(G)| = p^3$ in order that $|G/\Phi(G)| = p^2$. Hence, $Z(G) = \Phi(G)$.

Choose $a, b \in G$ such that $\langle a, b \rangle = G$. By the previous paragraph, $a^p, b^p \in Z(G)$ so $|a|, |b| \leq p^2$ since $Z(G)$ is elementary abelian.

To finish, $[a, b] \in Z(G)$ and it is easy to see any $x \in G$ has a representation $x = a^i b^j [a, b]^k$ for $i \leq |a|$, $j \leq |b|$, $k \leq |[a, b]|$. This is because a, b generate G , any instance of aba in the representation for an element can be replaced with $aab[a, b]$, and $[a, b] \in Z(G)$. Since $|G| = p^5$, and $|[a, b]| \leq p$ (as $Z(G)$ is elementary abelian), then $|a| = |b| = p^2$ and $|[a, b]| = p$.

Therefore, G has the same presentation as our C-group of the same size, so the two are isomorphic. $\square$

Fact 1. *All C-groups of the class defined above are nilpotent of class 2.*

Proof. Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$ in some C-group G of the class defined above. Then $[x, y] = (y_2 x_3 + (n_2 - 1)x_2 y_3, 0, 0)$. So $[x, y]$ is clearly in the center of G . Therefore, all C-groups in our class are nilpotent of class 2. $\square$

More C-Groups

The multiplication defined for α C-groups appears to be a semidirect product, but GAP tests indicated that was not the case. Though the number of α C-groups is infinite and can be extended infinitely by right direct product with an appropriate abelian group such as $\mathbf{Z}_{kn_1}$ or $\mathbf{Z}$ where k is a positive integer, this class does not describe all C-groups. For example, there is a C-group of order 64 that is nilpotent of class 3 with the following group representation

$$\langle a, b \mid a^4 = b^8 = 1, a^2 b = b a^2, b^2 a = a b^{-2}, (b^{-1} a)^2 = (a b)^2 \rangle.$$

There are also two C-groups of order 96 that are not nilpotent. These facts raise some interesting questions: Is there a bound on the nilpotency class of C-groups? What is the structure of the nilpotent C-groups not in our class? What is the structure of non-nilpotent C-groups? In addition, all C-groups of order less than 1000 are solvable. Are all C-groups solvable?

A brief examination of the GAP data reveals that the number of generators of the center of C-groups of order less than 1000 is a linear function of the number of group generators. We conjecture that for finite C-groups, the number of generators of the center is polynomially bounded by the number of generators of the group. Clearly, there is no such bound for C-groups with infinitely generated centers.

It is also interesting to note that for the order of any C-group less than 1000 there is a C-group in the class described above. Does this hold for orders greater

than 1000? That is, can we have a C-group of an order not covered by our class?

Kazem Mahdavi
Department of Mathematics
SUNY Potsdam
Potsdam NY 13676
mahdavr@potsdam.edu

Avi Silterra
5714A Beacon St
Pittsburgh PA 15217
ajs2@andrew.cmu.edu

Mihai

Margarethe

References

- [1] Abels, Herbert, "An example of a finitely presented solvable group," *London Mathematical Society lecture notes*, Cambridge: Cambridge University Press, 1971, pp.205-211.
- [2] Dummit, David S., Richard M. Foote, *Abstract Algebra*, Hoboken: John Wiley & Sons, Inc., 1999.
- [3] The GAP Group - Groups, Algorithms, and Programming, Version 4.3; 2002.
(<http://www.gap-system.org>)
- [4] Hungerford, Thomas W., *Algebra*, New York: Springer-Verlag, 1974.
- [5] Robinson, Derek John Scott, *A course in the theory of groups*, 2nd Edition, New York : Springer-Verlag, 1995.